

Appendix J

Fisheries Surveys



Fisheries Assessment for the N6 Galway City Transport Project

May 2018



Table of Contents

1.	Introduction	2
2.	Methodology	6
3.	Results	12
4.	Discussion	36
5.	References	41

List of Tables

- Table 1.1 - Watercourses surveyed as part of the proposed road development
Table 3.1 – Macro-invertebrate species composition of riverine sites downstream of the proposed road development
Table 3.2 – Summary of fisheries results for each riverine catchment surveyed for the proposed road development
Table 3.3 – Summary of fisheries results for each riverine catchment
Table 3.4 - Depletion statistics for the Trusky, the Bearna and Knocknacarragh Streams
Table 3.5 – Fish species recorded during fyke net surveys at Ballindooley Lough and Coolagh Lakes
Table 3.6 – Fisheries Evaluation of Watercourses surveyed as part of the proposed road development

List of Figures

- Figure 1.1 – Watercourses surveyed
Figure 2.1 – Location of Fisheries Survey Sites along the proposed road development
Figure 3.1 - Q sampling sites downstream of the proposed road development
Figure 3.2 - Biological Q-Ratings of riverine sites downstream of the proposed road development
Figure 3.3 - Length Frequency distribution of fish species captured in Sruthán na Líbeirtí
Figure 3.4 - Length Frequency distribution of fish species captured in the Trusky Stream
Figure 3.5 - Length Frequency distribution of fish species captured in the Bearna Stream
Figure 3.6 - Length Frequency distribution of fish species captured in the Tonabrocky Stream
Figure 3.7 - Length Frequency distribution of fish species captured in the Knocknacarragh Stream
Figure 3.8 - Length Frequency distribution of fish species captured in the Terryland River
Figure 3.9 - Depletion curve calculated for the Bearna Stream
Figure 3.10 - Depletion curve calculated for the Trusky Stream
Figure 3.11 - Depletion curve calculated for the Knocknacarragh Stream
Figure 3.12 – Length Frequency distribution of fish species captured at Ballindooley Lough
Figure 3.13 – Length Frequency distribution of fish species captured at Coolagh Lakes
Figure C1 - Location of Fyke Netting Sites on Ballindooley Lough
Figure C2 - Location of Fyke Netting Sites on the Coolagh Lakes

1. Introduction

Triturus Environmental Services were contracted by Scott Cawley Ltd. (ecological consultants) to conduct a fisheries assessment of watercourses along the corridor of the proposed N6 Galway City Transport Project, here and after referred to as the proposed road development. This included watercourses crossed by the proposed road and of watercourses with downstream hydrological connectivity. The work was prepared to collect baseline fisheries data that would inform the preparation of the Environmental Impact Assessment prepared as part of the planning requirements.

Triturus Environmental Services made an application under Section 14 of the Fisheries (Consolidation) Act, 1959 as substituted by Section 4 of the Fisheries (Amendment) Act, 1962, to undertake quantitative electro-fishing surveys of small rivers/ streams along the corridor of the proposed road alignment development (see Figures 3.1 and 3.3). Small river and stream habitats were thus surveyed by electro-fishing or if a very minor channel, appraised for their fisheries value. The surveys were conducted at intersections with the proposed road development and at accessible downstream locations. This included an assessment of each riverine habitat as spawning, nursery and holding habitat for fish.

The field survey was designed to help contextualise sensitive fisheries habitat relative to the proposed road development works, including the downstream habitat that may be impacted from water borne pollutants. Furthermore, baseline biological water quality (i.e. Q-values) were collected at each watercourse crossing where suitable habitat existed. This will also help relate water quality baselines to fish population data (i.e. clean water salmonid etc.). The biological water quality collected would also provide baseline readings against which future water quality targets could be gauged. These values should not deteriorate as a result of works associated with the proposed road development. According to the Water Framework Directive (2000/60/EEC), all watercourses should aim to maintain or achieve target 'good status' water quality.

Background

The proposed road development is approximately 17km in length. As described moving from the west eastwards, the proposed road development ties into the existing R336 Coast Road at an at-grade roundabout junction (Bearna West Roundabout) approximately 2km to the west of Bearna Village (see Figure 1.1). The proposed road development then proceeds north before veering eastwards to the north of Bearna Village and onwards towards the townland of Letteragh to a grade separated junction (N59 Letteragh Junction) to connect the N59. The N59 Link Road connects the proposed road development to the N59 to the north of Bushypark Church and to the Letteragh Road and Ragoon Road to the south.

The proposed road development then continues in sections of cut and fill, travelling over the N59 at Dangan, through National University of Ireland Galway's recreational lands and over the River Corrib on a bridge structure. To the east of the river, the proposed road development is generally on embankments or viaduct structures, before entering a section of cut preceding a tunnel at Lackagh Quarry. After exiting the tunnel, it continues on embankments and passes over the N84. Grade separated signalised junctions are provided at the N84 (N84 Headford Road Junction) and the N17 (N17 Tuam Road and Parkmore Junction). The proposed road development then proceeds in a cut and cover tunnel to the north of Galway Racecourse before turning south to connect to a free-flow grade separated junction to the south of the existing Briarhill Junction, in the townland of Coolagh, Briarhill. The proposed road development then ties in to the existing N6 dual carriageway at Coolagh, Briarhill.

The proposed road development overlaps three hydrometric areas (HA 29, 30 & 31) containing the watercourses that formed the basis of this fisheries assessment (see Figure 1.1 below). The watercourses overlapping or connected downstream of the proposed road development were dominated by small to medium stream and river habitats that included Sruthán na Libeirtí, Trusky Stream, Bearna Stream & tributary, Tonabrocky Stream, Knocknacarragh Stream, Terryland River and Merlin Stream (see Table 1.1 below). The larger rivers and stream sites were visible on the OSI 1:50,000 scale mapping. On the OSI mapping rivers were defined as dark blue lines and streams defined as light blue lines. Smaller streams or ephemeral (seasonal) streams and ditches were not visible. In addition to the main channels of the streams listed above, small low order tributaries that were not consistently visible on OSI 1:50,000 scale mapping, were also appraised. These small watercourses were connected hydraulically to the larger stream catchments and were identified from hydrological GIS map layers. Two lake systems were also surveyed given downstream hydraulic connectivity to the proposed road development. These were the Coolagh Lakes and Ballindooley Lough (immediately north of Galway City) both of which are contained within the Lower Corrib catchment and within hydrometric area 30. Ballindooley Lough is an isolated valley basin lake and not connected to a major river system. In contrast the Coolagh Lakes are connected to the River Corrib by a small reeded channel. All of the watercourses within the study site are located within the Western River Basin District.

The streams west of the River Corrib, and contained within hydrometric area 31, are situated on geologies of Siluro-Devonian Granitic Rocks and Appinite according to the Geological Survey of Ireland. These watercourses may be considered less alkaline as a result of the underlying geologies, as was reflected by the plant communities present including alternate water milfoil *Myriophyllum alterniflorum* and *Fontinalis squamosa*.

The watercourses east of the River Corrib (i.e. Terryland River, Merlin Stream, Coolagh Lakes and Ballindooley Lough) are situated on Visean Limestone and Calcareous Shale. These watercourses are therefore considered more alkaline and this was reflected by the macrophyte

plant communities present that included common club rush *Schoenoplectus lacustris*, Chara species and Alkaline fen vegetation (i.e. Ballindooley Lough).

Table 1.1. Watercourses surveyed as part of the proposed road development

Watercourse name	Type	EPA Code	Hydrometric Area	River Basin
Sruthán na Libeirtí	Medium sized stream	31F01	31	Western
Trusky Stream	Medium sized stream	31B02	31	Western
Bearna Stream & tributary	Large Stream	31O05	31	Western
Tonabrocky Stream (forms tributary with the Bearna Stream at Ballard)	Large stream	31B01	31	Western
Knocknacarragh	Small stream	31K16	31	Western
Terryland River	Large stream	30T01	30	Western
Merlin Stream	Seasonal stream	No EPA code	29	Western
Coolagh Lakes & river tributary	Natural lake	No EPA code	30	Western
Ballindooley Lough	Valley Fen Lake	No EPA code	30	Western




Figure 1.1. Watercourses surveyed (survey sites highlighted in purple with proposed road development highlighted in red) prepared using Quantum GIS 2.18)

2. Methodology

This section summarises the methodologies employed for the various sub components of the site surveys. It includes the following elements as described in the sections below, macro-invertebrate (Q sampling), electro-fishing practice, lamprey surveys, salmonid surveys, optimum survey times and bio-security protocol.

Macro-invertebrate (Q Sampling)

Macro-invertebrate samples were collected at the stream sites crossed by the proposed road development between the 22nd and 30th September 2015 in advance of the fisheries surveys. Samples were collected at the nearest location containing riffle/ glide habitat downstream of proposed crossings. The samples were collected by 'kick' sampling for approximately 2.5 minutes in the faster flowing areas (riffles) of the streams using a standard hand net (250 mm width, mesh size 500 micron). The samples were collected from seven stream sites, Sruthán na Libeirtí, Trusky Stream, Bearna Stream tributary, Bearna Stream, Tonabrocky Stream, Knocknacarragh Stream and the Terryland River (see Figure 3.1 below for locations). The samples were collected by moving across the riffle zone and also involved washing large rocks from the riffle zone to ensure a full representation of the species composition. Collected samples were elutriated, and fixed in 70% ethanol prior to identification.

The macro-invertebrates were later identified using a Nikon SMZ 1000 stereo microscope and Freshwater Biological Association invertebrate keys. Invertebrate taxa were identified to species level where possible and grouped based on the Environmental Protection Agency (EPA) categories from pollution intolerant to very pollution tolerant on a moving scale from A to E (see Appendix I of Water Quality in Ireland 2001-2003, Toner et al., 2005).



Plate 1.1 – Nikon SMZ 1000 microscope and FBA keys used for macro-invertebrate identification

Electro-fishing practice

An electro-fishing survey of the existing fish stocks within each watercourse was undertaken between the 22nd of September and 30th September 2015 in the environs of Galway City (see Figure 2.1 below for site locations).

The surveys were undertaken along sections of watercourses crossed by the proposed road development or, where the channel was seasonal or inaccessible, at the closest location downstream. The survey sections were 50m in length and sealed off with stop nets, effectively acting as fish barriers for the depletion survey.

Water with a low conductivity has a higher resistance to the passage of an electric current through it. This means that in high conductivity waters the current for a given voltage is higher than in low conductivity water and the threshold values for different fish responses are also lower (Zalewski and Cowx, 1990). Given this fact, the lower conductivity waters to the west of the River Corrib (i.e. Sruthán na Libeirtí, Trusky and



Plate 1.2 – Electro-fishing the Terryland River

Tonabrocky) were fished between 250-300 volts salmonids / 100volts for lamprey. In the more alkaline watercourses to the east of the River Corrib electro-fishing was conducted at 225volts for salmonids / 100volts for lamprey. Similarly where no suitable upstream habitat was available (as was the case at two sites (i.e. Trusky and Knocknacarragh Streams) the high conductivity transitional reaches were surveyed (i.e. adjoining tidal channel sections in the lower Trusky & and Knocknacarragh streams). Consequentially a lower voltage of 75-100volts was utilised given the higher conductivity. Other settings i.e. frequency, duty cycle etc. are discussed below. Depletion electro-fishing of each site was conducted by two operators in an upstream direction using a single anode Smith-Root LR24 backpack (12V DC input; 300V, 100W DC output). In order to minimise potential damage and undue stress to lamprey species and Atlantic salmon, electro-fishing settings were modified to target specific species at the site (see lamprey section below). Larval lamprey species, for example, were specifically targeted in areas of low/reduced flow and with a higher proportion of soft sediment. However, this habitat was recorded as very localised or entirely absent in many watercourses with the exception of the Terryland River.

Salmonids typically require a higher frequency (and also voltage) than lamprey species in order to sufficiently stun them for capture. Unless amended, these settings can result in the inadvertent electro-narcosis of buried ammocoetes, resulting in failure to emerge and recording of absence, as well as damage to the fish (Thompson et al. 2010). Therefore soft sediment areas were identified and targeted first following stop netting.

Fisheries Habitat Suitability

Small stream habitats (typically less than 1m wide) were assessed for their fisheries habitat suitability (see Figure 2.1 below). These streams were typically heavily choked with vegetation or seasonal and in general were considered unsuitable for electro-fishing.

These were characteristically degraded watercourses of low or no fisheries value. Nonetheless they were still visited to validate whether any fisheries value existed. The appraisal included for spawning habitat (i.e. fast water and clean gravel habitat for recruiting adults), nursery habitat (i.e. juvenile fish habitat with faster riffle habitat for salmonids and good instream structure or fine sediment for lamprey) and holding habitat (deeper pool habitat with cover for adult fish).

Lamprey

Settings for lamprey followed those recommended and used by APEM (2004) and Niven & McAuley (2013). Using this approach, the anode was placed under the water surface, approximately 10–15 cm above the sediment, to prevent immobilising lamprey ammocoetes within the sediment. The anode was energised with 100V of pulsed DC for 15-20 seconds and then turned off for approximately five seconds to allow ammocoetes to emerge from their burrows. The anode was switched on and off in this way for approximately two minutes. Immobilised ammocoetes¹ are collected by a second operator using a fine-mesh hand net as they emerge. During this survey, settings utilised to capture juvenile lamprey were a frequency of 25Hz, voltage of 100V and a duty cycle of approximately 12% (pulse width 6ms).

Salmonids

As salmonids typically require a higher frequency than lamprey ammocoetes, the frequency was set at 40Hz frequency, a voltage of between 225V and 300V (dependant on local water conductivity) and a duty cycle of 18%. The higher voltage was used in waters west of the River Corrib, where conductivity was lower.

Multiple-pass depletion electro-fishing methodology was employed and followed those outlined by Carle & Strub (1978) and Lockwood & Schneider (2000). Population estimates were made utilising the following equations:

$$T = \sum_{i=1}^k C_i,$$
$$X = \sum_{i=1}^k (k-i)C_i,$$

¹ Ammocoetes are the juvenile larval stage of lamprey

where,

i = pass number

k = number of removals (passes)

C_i = number of fish caught in the sample

X = an intermediate statistic used below

T = total number of fish caught in all passes

The maximum likelihood estimate of N was determined by an iterative process by substituting values for n until:

$$\left[\frac{n+1}{n-T+1} \right] \prod_{i=1}^k \left[\frac{kn-X-T+1+(k-i)}{kn-X+2+(k-i)} \right] \leq 1.0,$$

Where n is the smallest integer satisfying the above equation.

The probability of capture, p , and variance of N are then estimated by:

$$p = \frac{T}{kN - X},$$

$$\text{Variance of } N = \frac{N(N-T)T}{T^2 - N(N-T) \left[\frac{(kp)^2}{(1-p)} \right]},$$

$$\text{Standard error of } N = \sqrt{\text{Variance of } N}.$$

Additionally, a goodness of fit test evaluating the power of capture probability was also implemented following the method of White et al. (1982) where;

$$E(C_1) = Np,$$

and for $i > 1$.

$$E(C_i) = N(1-p)^{i-1} p$$

Calculated χ^2 then is:

$$\chi^2 = \frac{[C_1 - E(C_1)]^2}{E(C_1)} + \frac{[C_2 - E(C_2)]^2}{E(C_2)} + \dots + \frac{[C_k - E(C_k)]^2}{E(C_k)}.$$

Fyke Netting

Boat based fyke netting surveys were undertaken at Ballindoooley Lough and Coolagh Lakes (see Figure 2.1 below for site locations). In advance of setting the nets a high resolution transducer was used to locate fish markings and establish a depth profile of the lake basins. This facilitated the positioning of the fyke nets near shelf drop offs and helped establish distributional patterns of fish. Five 1.5 meter diameter (D shaped) fyke nets with multi panel mesh were placed in the margins of the lakes in the littoral zones (windward bank) and shallow bay areas overnight and retrieved within 24 hours. The fish captured were measured by two personnel and length frequency graphs and species composition graphs were constructed. All fish were processed quickly and returned alive to the lakes.

Optimum Survey Period

The electro-fishing survey was undertaken between the 22nd and 30th of September 2015. This choice of survey period avoided a clash in spawning and upstream migration of both lamprey species and Atlantic salmon. It is considered that by undertaking electro-fishing surveys for salmonids during or post-August (in this case September) that the juvenile fish (including young-of-the-year individuals) are of sufficient size to recover following a survey which was conducted according to Inland Fisheries Ireland best practice (IFI pers. comm. 2015). In addition, the metamorphosis of lamprey usually takes place between July and September. It is therefore recommended that surveys for ammocoetes are carried out in July at the earliest but preferably between August and October in order to detect the presence of transformed ammocoetes (Harvey & Cowx, 2003). Additionally, by undertaking fyke netting before the end of September, it facilitates the detection of fish species that during the colder periods (i.e. winter) move to the deeper water of lakes and thus can remain undetected.

Biosecurity Protocol

All equipment and personal protection equipment (PPE) used was disinfected with Virkon® prior to and post-survey completion. Best practice precautions were employed to prevent the potential spread of invasive species and water-borne pathogens, according to standard Inland Fisheries Ireland (IFI) biosecurity protocols.




Figure 2.1. Location of Fisheries Survey Sites along the proposed road development (prepared using Quantum GIS 2.18)

3. Results

A) Biological Water Quality

Existing Water Quality Records

No biological water quality sampling has been previously recorded by the EPA in the stream sites overlapping the proposed road development. As such biological water data was collected immediately downstream of the proposed road crossing locations (where feasible) during September 2015 to determine baseline biological water quality for each of the stream sites surveyed. Where the channel was dry, too deep and thus unsuitable for Q sampling the closest downstream location was chosen to collect the sample. The biological water quality data recorded during the surveys is summarised below.

The EPA Q Sampling methodology is described in the EPA document Water Quality in Ireland 2001-2003 (Toner et al. 2005). The system groups macro-invertebrates into classes (A-E), whereby pollution intolerant species are denoted class A and species with greater pollution tolerances fall into successive classes (i.e. class E being very pollution tolerant). As such the presence or absence of these groups and their relative abundances facilitates an assessment of biological river health known as a Q Rating with Q1 being the poorest water quality rating and Q5 the highest water quality rating. The results of the 2015 data collection are discussed in this context in order to interpret changes in the river community composition. Refer to Figure 3.1 below for locations and Figure 3.2 for a summary of the Q Ratings at each of the sites surveyed (i.e. sites 1 through 7 downstream of the proposed road development). Table 3.3 lists all of the macro-invertebrate species recorded and represents by colour separation the EPA taxonomic classes as prescribed above. A summary of the Q Ratings are also provided in the table below.

Site 1 was located on Sruthán na Libeirtí, a small upland eroding watercourse heavily encroached by bracken, bramble and gorse scrub. The stream site evidently was suffering from organic enrichment given the exuberant growth of lesser water parsnip *Berula erecta*, fool's watercress *Apium nodiflorum* and common duckweed *Lemna minor* where light penetration permitted growth. This was also reflected in the macro-invertebrate community by the absence of cleanwater stoneflies and mayflies. The dominance of crustaceans (*Aselus aquaticus* and *Gammarus duebenii*), particularly the pollution tolerant *Aselus aquaticus* in very high numbers indicates at least moderate levels of pollution. The absence of both class A and B clean water taxa accounted for a Q rating of 3 in An Sruthán na Libeirtí.

The Trusky Stream (main channel) rises at An Chloch Scoilte north of Bearna. The stream then flows west and south adjoining two small low order tributaries in Bearna Village before discharging to the sea adjacent to Bearna Pier. The Trusky stream catchment, comprised predominantly shallow drainage channels that were heavily vegetated with herbaceous species including yellow flag *Iris pseudacorus*, fool's watercress, lesser water parsnip and common starwort *Calitriche stagnalis* instream.

The upper reaches of the catchment channels were largely dry with the exception being the tributary at An Chloch Scoilte north of Bearna Village that had shallow flowing water. The kick sample collected here on the Trusky Stream had a very low diversity of macro-invertebrates (n=5), possibly as a consequence of being seasonal (likely dries up in warm years) and also as a result of evident organic enrichment. The species composition was dominated by *Aselus aquaticus* and *Gammarus duebenii* and had no clean water stoneflies, mayflies or cased caddis species present. As such a Q rating of 3 was recorded in the Trusky Stream.

A small tributary of the Bearna Stream rising in the valley escarpments of 'Na hAille' to the west formed a confluence with the Bearna Stream north of Ballard. The stream was located in very dense bracken and gorse scrub and was shallow and swift flowing. The stream substrata comprised a coarse gravel and cobble base. It was seasonal further downstream, grading into a vegetated ditch on its approach to the Bearna Stream from the west. The stream had a low species diversity (n=5) dominated by casless caddis (*Plectonemia conspersa* & *Wormaldia occipitalis*) and pollution tolerant gastropods species (*Lymnaea peregra* & *Planorbis carinatus*). Given the seasonal nature of the stream and the dominance of class C and D invertebrates, a Q value of 3 was recorded at the site.

The Tonabrocky Stream rises at Tonabrocky and flows south west into Rusheen Bay, west of Rusheen Point. It has a large tributary, the Bearna Stream, that rises at Lough Cnoc an Champa and then flows south east before forming a confluence with the Tonabrocky Stream at Ballard. The Tonabrocky Stream and its tributary the Bearna Stream are both fast flowing salmonid streams with clean gravels and limited macrophyte plant growth with the exception of liverworts, mosses and small stands of alternate water milfoil *Myriophyllum alterniflorum*. Localised sections also have stands of lesser water parsnip and fool's watercress. The evident low levels of enrichment were reflected in the macro-invertebrate communities of both streams that contained three families (Chloroperlidae, Nemouridae & Leuctridae) of clean water stonefly species (EPA Class A and B). The streams also contained clean water cased caddis families (EPA class B) Odontoceridae and Seracostomatidae. The presence of good numbers of EPA class A and B macro-invertebrates indicated unpolluted water (Q4) in both the Tonabrocky Stream and its major tributary the Bearna Stream.

The Knocknacarragh Stream rises west of Letteragh and flows south west into Rusheen Bay (entering north of Blake's Hill). The upper catchment low order tributaries were predominantly seasonal drainage channels. However, one small tributary north-west of Ballyburke contained a shallow flowing stream where it was possible to collect a kick sample. The Knocknacarragh Stream itself is considered seasonal and this was reflected by the poor diversity of macro-invertebrates (n=6). It is situated in a catchment supporting beef cattle and was suffering from organic enrichment given the presence of abundant algae. This was also reflected by virtue of the absence of clean water macro-invertebrates with the exception of the (EPA class B) cased caddis species *Seracostoma personatum*.

This cased caddis species however is considered more tolerant to pollution than other cased caddis families (e.g. Goeridae, Limnephilidae, Odontoceridae etc.). The macro-invertebrates recorded at the site indicated an equivalent Q rating of 3 (i.e. moderately polluted water quality).

The Terryland River rises in the urbanised area of north east Galway City (i.e. Terryland). The river is situated on karstic limestone and unusually can flow in both directions, either towards the River Corrib at Jordan's Island or in the opposite direction towards Terryland. It was the largest of the river and stream sites surveyed. The presence of the nitrate loving *Sparganium erectum* in abundance, a species favouring soft bottomed enriched and modified channels indicated historical deepening/ straightening of the the river channel. Furthermore, enrichment of the stream was evident by the presence of the exuberant growth of water milfoil *Myriophyllum spicatum* and common starwort *Callitriche stagnalis* but also pondweeds *Potamogeton spp.* The macro-invertebrate community also indicated gross pollution. The absence of cleanwater Class A & B macro-invertebrates and the presence of pollution tolerant forms (EPA class E, *Chironomus riparius* & *Tubifex sp.*) indicated more heavily polluted water quality (i.e. Q2-3).

In summary the water quality was of poor status (i.e. Q2-3 & Q3) at all stream sites with the exception being the Bearna Stream and the Tonabrocky Stream where water quality was of good status (i.e. Q4). Figure 3.2 below presents the Q sampling results at each survey site. The red colour on the graph identifies Q3 moderately polluted water quality, orange colour identifies Q2-3 seriously polluted water quality and green colour identifies Q4 good water quality which is the Water Framework Directive target status.

Site 1 (Sruthán na Libeirtí)	Q3 - Moderately Polluted (Poor Status)
Site 2 (Trusky Stream)	Q3 - Moderately Polluted (Poor Status)
Site 3 (Bearna Stream tributary)	Q3 - Moderately Polluted (Poor Status)
Site 4 (Bearna Stream)	Q4 - Unpolluted (Good Status)
Site 5 (Tonabrocky Stream)	Q4 - Unpolluted (Good Status)
Site 6 (Knocknacarragh Stream)	Q3 - Moderately Polluted (Poor Status)
Site 7 (Terryland River)	Q2-3 - Seriously Polluted (Poor Status)




Figure 3.1. Q sampling sites downstream of the proposed road development (prepared using QGIS 2.18)




Figure 3.2. Biological Q-Ratings of riverine sites downstream of the proposed road development

Table 3.2. Macro-invertebrate species composition of riverine sites downstream of the proposed road development

Group	Family	Species	Site 1 Sruthán na Libeirtí Stream	Site 2 Trusky Stream	Site 3 Bearna Stream trib.	Site 4 Bearna Stream	Site 5 Tonabrocky Stream	Site 6 Knocknac arragh Stream	Site 7 Terryland River	EPA Class
Stoneflies (Class A)	Chloroperlidae	<i>Chloroperla torrentium</i>					9			A
		<i>Chloroperla tripunctata</i>				9	11			A
	Nemouridae	<i>Amphinemura sulcicollis</i>				3				A
Stoneflies (Class B)	Leuctridae	<i>Leuctra inermis</i>				6	5			B
Cased Caddis (Class B)	Seracostomatidae	<i>Sercacotoma personatum</i>				5	3	1		B
	Odontoceridae	<i>Odontocerum albicorne</i>				2				B
Mayflies (Class C)	Baetidae	<i>Baetis rhodani</i>				5	8			C
Caseless Caddis (Class C)	Hydropsychidae	<i>Hydropsyche siltalai</i>				1	1			C
	Polycentropdidae	<i>Plectronemia conspersa</i>	12		6				2	C
		<i>Polycentropus kingi</i>				1		1		C
	Philopotamidae	<i>Wormaldia occipitalis</i>			2					C
	Ryacophilidae	<i>Ryacophila dorsalis</i>					1			C
Crustaceans (Class C)	Gammaridae	<i>Gammarus duebenii</i>	47	29	4	7	23	8	76	C
Beetles (Class C)	Elmidae	<i>Limnius volckmari</i>	2			4	8	1		C
		<i>Elmis aenea</i>	9			2	1	5		C
		<i>Oulimnius sp.</i>				3	2			C
	Dytiscidae	<i>Dytiscus sp.</i>							2	C
	Hydraenidae	<i>Hydraena sp.</i>	5							C
Dipterans (Class C)	Chironomidae	<i>Orthocladius sp.</i>			2	2	3			C
	Simuliidae	<i>Simulium sp.</i>	6	4						C

Group	Family	Species	Site 1 Sruthán na Libeirtí Stream	Site 2 Trusky Stream	Site 3 Bearna Stream trib.	Site 4 Bearna Stream	Site 5 Tonabrocky Stream	Site 6 Knocknac arragh Stream	Site 7 Terryland River	EPA Class
	Tipulidae	<i>Tipula sp.</i>								C
Gastropods (Class C)	Hydrobiidae	<i>Potamopyrgus antipodarum</i>	5	31	11	4		6		C
	Succineidae	<i>Succinea putris</i>	1							C
	Valvatidae	<i>Valvata piscinalis</i>	1							C
		<i>Valvata cristata</i>							1	C
	Bithyniidae	<i>Bithynia tentaculata</i>							1	C
Freshwater Limpet (Class C)	Ancylidae	<i>Ancylus fluviatilis</i>	2	2	5	4				C
Freshwater Bug (Class C)	Corixidae	<i>Hesporixia linnaei</i>							4	C
	Lymnaeidae	<i>Lymnaea peregra</i>			2				2	D
	Planorbidae	<i>Planorbis carinatus</i>			1					No Class
Freshwater Hoglouse (Class D)	Aselidae	<i>Asellus aquaticus</i>	61	43	6		1		35	D
Leeches (Class D)	Erpobdellidae	<i>Erpodella octoculata</i>			1		1			D
	Glossiphoniidae	<i>Glossiphonia heteroclita</i>			1					D
Freshwater Worms (No Class)	Oligochaeta	<i>Stylodrilus heringianus</i>	3		1					No Class
Non-biting midge (Class E)	Chironomidae	<i>Chironomus riparius</i>							15	E
Freshwater Worms (Class E)	Tubificidae	<i>Tubifex sp.</i>							4	E
Total Abundance			153	109	52	54	64	22	148	
N			12	5	13	15	13	6	10	
Q Rating			Q3	Q2-3	Q3	Q4	Q4	Q3	Q2-3	

B) Fisheries

Stream Sites

An electro-fishing survey of the existing fish stocks on stream sites overlapping the corridor of the proposed road development was conducted between the 22nd of September and 30th September 2015, following notification to Pat Gorman, regional inspector for Inland Fisheries Ireland Galway. The results of the survey are discussed below in terms of fish population structure and the typical value of the surveyed areas as nursery, spawning and holding habitats for various fish species. Seven stream catchments were electro-fished (refer to Table 3.2 for a list of the surveyed watercourses and Figure 2.1 for a location map). These were typically small streams between 0.5m and 2m in width, some of which were seasonal and not of fisheries value. The Terryland River was the largest watercourse surveyed at between 5-8m in width. One large stream channel, the Merlin Stream was surveyed at two sites but did not contain water at either site at the time of the survey and thus is not discussed further. A summary of the recorded fish species and their abundances is provided in Table 3.2. Length-frequency plots of fish species recorded at each stream site are provided in Figures 3.4-3.9 and described in the text below. Depletion curves for electro-fishing effort are illustrated in Figures 3.10-3.12 and also described in the text below.

Sruthán na Libeirtí

No fish were captured from site (1A) in the upper reaches of the Sruthán na Libeirtí Stream. While some apparent moderate quality salmonid habitat existed (i.e. riffle, glide and pool habitat) with cobble and gravel substrata no fish were recorded as present. Records show that the stream dried up in the upper reaches during the warm summers of 2013-2014 and this may account for the absence of fish. The stream has also been modified historically (localised straightening & deepening). Further downstream at site 1B (circa 200m from the sea), two European eel *Anguilla anguilla* were captured. Both were silvered and ready to return to sea to spawn (see Plate 3.1 and Figure 3.3 below). The lower reaches of the Sruthán na Libeirtí stream had good flows of water, averaging 0.4m deep with good quantities of gravel and cobble. Again while some moderate quality salmonid habitat existed no salmonids were recorded during the survey.



Plate 3.1. Sruthán na Libeirtí

Trusky Stream

Two sites were surveyed on the Trusky Stream. Much of the upper catchment and lower order tributaries were seasonal ditches that were heavily vegetated channels, with the exception of a small flowing stream at An Chloch Scoilte (site 2A). Here the stream had a cobble and gravel base and a good profile with riffle and glide habitat and localised pool. Despite an electro-fishing effort over two 25m sections no fish were recorded as present.

The compacted and bedded gravels in addition to the shallow nature of the stream indicated it would have limited capacity to support salmonids. The stream was also suffering from organic enrichment from agriculture in the upper catchment. As such, a secondary site (2B) was surveyed downstream at Bearna Village, where greater flows of water and a larger channel were present given that the site was below the confluence of three smaller tributaries. This section of channel was tidal (on spring tides) and proved to be an excellent nursery for flounder *Platichthys flesus*, European eel and three-spined stickleback *Gasterosteus aculeatus* (see Figure 3.4). Small numbers of adult brown trout *Salmo trutta* were also present. The brown trout population was small with only two fish present and given the limited better quality habitat available upstream the total stream population size must also be comparably small. Some moderate to good quality spawning habitat did exist in the lower 100m of the stream catchment. This area is likely to be the main area for recruitment in the stream given the poor quality upstream salmonid habitat.



Plate 3.2. Trusky Stream

Bearna Stream

The Bearna Stream was an excellent salmonid river with widespread clean and un-compacted spawning gravels and swift flows of water in the corridor for the proposed road development (Site 3B). The river had a well defined profile, characteristic of good quality salmonid habitat with stretches of riffle grading into glide and pool habitat. This created an excellent salmonid nursery habitat and good areas of adjoining pool to support older adult year classes. This was reflected in the good numbers of brown trout recorded at the site (see Figure 3.5). European eel were also recorded present further exemplifying the high quality fisheries value of the stream. A small tributary of the Bearna Stream was also surveyed (i.e. site 3A) but given it was a shallow, small (<0.5m wide) and seasonal stream no fish were recorded present.



Plate 3.3. The Bearna Stream

Tonabrocky Stream

The Tonabrocky Stream in its upper reaches in the vicinity of the proposed road development had poor quality fisheries habitat being predominantly a dry drainage channel with localised pockets of water. Downstream however the habitat improved significantly, turning into a swift flowing stream with a gravel base and clean water (e.g. site 4A). Here small numbers of brown trout were recorded (see Figure 3.6). At this location and all the way downstream to Rusheen Bay, the stream had good quality salmonid habitat with nursery, spawning and holding habitat present over much of the channel length.



Plate 3.4. Tonabrocky Stream

Knocknacarragh Stream

The upper reaches of the Knocknacarragh Stream were of poor fisheries value (e.g. site 5A). No fish were recorded in the upper reaches. However, a small stretch of water emanating from a culvert on approach to Rusheen Bay (site 5B) proved to be a good nursery for estuarine fish and small numbers of European eel. At this location the stream became tidal with a cobble, sand and gravel base that supported sand goby *Pomatoschistus minutes*, grey mullet *Chelon labrosus*, three spined stickleback and small numbers of European eel (see Figure 3.7).



Plate 3.5. Knocknacarragh Stream

Coolagh Lakes tributary

A small tributary entering the western bank of the upper Coolagh Lakes (site 6A) was surveyed to establish whether the habitat was of value to salmonids, eel or lamprey. The channel was deep 1.2-2.0m and was heavily vegetated with lesser water parsnip, fool's watercress and common reed *Phragmites australis* and reed canary grass *Phalaris arundinacea*.



Plate 3.6. Coolagh Lakes tributary

Despite electro-fishing two 25m sections in sequence no fish were recorded as present. The channel was largely stagnant and had a soft base with limited gravels. It therefore was not considered of importance to salmonids and suboptimal for lamprey. The channel most likely is of some value to coarse fish during the spring when species such as roach present in Coolagh Lakes may migrate to the vegetated channel to spawn.

Terryland River

The Terryland River was surveyed at two locations (7A & 7B), with only two European eel being captured during the survey at the lower site (7B) (see Figure 3.8). The river site was predominantly deep (0.9-2.2m) and heavily vegetated with pondweeds, water starwort, strap weeds and spiked water milfoil. The watercourse was evidently heavily enriched from urban storm water runoff and other sources. It also appeared to have been straightened and deepened historically. As such very little natural habitat remained and as a consequence was of limited fisheries value apart from European eel that can access the channel via its connection with the River Corrib at Jordan's Island.



Plate 3.7. Terryland River

Table 3.3. Summary of fisheries results for each riverine catchment surveyed

River Catchment	No. E-Fish Sites	Site No.	No. Fish Species Recorded (species in parenthesis)	Total number of fish (n)	Target Notes
1 - Sruthán na Libeirtí Stream	2	1A	None	0	Seasonal stream with some semi natural salmonid and eel habitat. Stream suffering from organic enrichment and has had historical channel modifications.
		1B	1 (European eel)	2	Moderate quality eel and salmonid habitat, however salmonids recorded as absent.
2- Trusky Stream	2	2A	None	0	Seasonal stream with some semi natural salmonid habitat. Stream has been impacted by organic enrichment.
		2B	4 (Flounder, European eel, 3 Spined Stickleback, Brown trout)	43	Swift flowing and good quality nursery and spawning habitat in lower reaches (approximately 100m of better quality habitat, upper system poor).
3A – Bearna Stream	2	3A	None	0	Seasonal stream with some semi natural salmonid and eel habitat.

River Catchment	No. E-Fish Sites	Site No.	No. Fish Species Recorded (species in parenthesis)	Total number of fish (n)	Target Notes
					Further downstream fisheries habitat poor as stream grades into seasonal ditch and wet grassland before forming confluence with the Bearna Stream.
		3B	2 (Brown Trout, European eel)	32	Excellent quality salmonid and eel habitat. Very good nursery and spawning with moderate quality adult holding habitat.
4 – Tonabrocky Stream	1	4A	1 (Brown Trout)	2	Moderate quality trout stream with low density of juvenile trout, perhaps colonising from better quality downstream habitat. Further upstream channel becomes a seasonal drainage channel.
5 – Knocknacarragh Stream	2	5A	None	0	Most western tributary in Knocknacarragh catchment has some moderate quality salmonid habitat but is seasonal. No fish recorded during survey. Other upper tributaries heavily modified and generally shallow, without water or culverted (moving downstream).
		5B	4 (European eel, Grey mullet, Sand goby, Flounder)	48	Transitional habitat emerging from culvert in lower reaches on approach to Rusheen Bay a good nursery for estuarine fish and to a lesser extent European eel despite evident sewage waste water present.
6 – Coolagh Lakes Stream tributary	1	6A	None	0	Deep slow moving section of drainage ditch connected to the upper Coolagh Lake. May be of importance for spawning coarse fish species during spring but not considered of value to salmonids or lamprey. None recorded during survey.
7 – Terryland River	2	7A	None	0	Both sites on the Terryland River were suffering from serious pollution (gross organic enrichment and other sources). Exuberant plant growth and historical channel deepening and straightening. Only 2 adult eel recorded during survey.
		7B	1 (European eel)	2	




Figure 3.3. Length Frequency distribution of fish species captured in Sruthán na Líbeirtí Stream




Figure 3.4. Length Frequency distribution of fish species captured in the Trusky Stream




Figure 3.5. Length Frequency distribution of fish species captured in the Bearna Stream




Figure 3.6. Length Frequency distribution of fish species captured in the Tonabrocky Stream




Figure 3.7. Length Frequency distribution of fish species captured in the Knocknacarragh Stream




Figure 3.8. Length Frequency distribution of fish species captured in the Terryland River

Fish population size estimation

Fish population size estimates were calculated using the stream fish population estimate methodologies of Carle & Strub (1978) and Lockwood & Schneider (2000). The population estimates were calculated for river sites where >20 fish were recorded. Smaller numbers of fish do not fit the Carl & Strub equations. As such, the depletion statistics were only calculated for the Trusky, the Bearna and Knocknacarragh Streams, as sufficient quantities of fish were captured, in these watercourses.

The accuracy of the estimated N was tested using the goodness of fit test (as per White et al., 1982). Given that $\chi^2 < \chi_{0.95}^2$ for the three watercourses tested there was no significant difference between the calculated capture probabilities, meaning that the rate of fish removal (depletion) can be considered as constant and in agreement with the depletion curves calculated (see Figures 3.10-3.12 and Table 3.3 below). The consistent depletion was achieved given experienced operators of electro-fishing equipment and the efficient use of stop nets. The correct use of the Smyth-Root electro-fishing gear current settings in a range of site conditions with different levels of capture efficiency (i.e. high conductivity, low conductivity, cobble / boulder dominated, deep, shallow, weeded etc.) ensured that population estimates could be improved.

Table 3.4. Depletion statistics for the Trusky, Bearna and Knocknacarragh Streams

Site	Captured population	Carl & Strub Pop. Est. with 95% confidence limits (NL & NU) in parenthesis	χ^2 (Goodness of fit)	$\chi^2 < \chi_{0.95}^2$	Constant Rate Depletion
Trusky	43	37 (46, 55)	1.276	(5.991, d.f.=2, p=0.581)	Yes
Bearna	31	36 (33, 37)	1.090	(5.991, d.f.=2, p=0.574)	Yes
Knocknacarragh	48	61 (35, 87)	0.817	(5.991, d.f.=2, p=0.333)	Yes




Figure 3.9. Depletion curve calculated for the Bearna Stream




Figure 3.10. Depletion curve calculated for the Trusky Stream




Figure 3.11. Depletion curve calculated for the Knocknacarragh Stream



Plate 3.8. Silvered European eel (*Anguilla anguilla*) captured in the lower reaches of the Sruthán na Libeirtí Stream



Plate 3.9. European eel elvers captured in the lower reaches of the Trusky Stream



Plate 3.10. Brown trout (*Salmo trutta*) captured in the Trusky Stream



Plate 3.11. Young Brown trout captured in the salmonid nursery of the Bearna Stream



Plate 3.12. Juvenile flounder (*Platichthys flesus*) captured in the transitional reaches of the Knocknacarragh Stream



Plate 3.13. Three spined stickleback (*Gasterosteus aculeatus*) captured in the Trusky Stream

Lake Sites

Ballindooley Lough and Coolagh Lakes (map sites 8 & 9 respectively on Figure 2.1) were surveyed between the 22nd and 24th of September 2015. The nets were positioned to maximise the capture rates from the lake meso-habitats (see Appendix B for depth profiles and Appendix C for fyke net locations). The positioning of the fyke nets included the windward bank on the Chara zones of the shelving margins, shallow bays and at oblique angles to natural points in the reed swamp littoral zones.

A total of four fish species were recorded from Ballindooley Lough. These included benthivorous tench *Tinca tinca*, pelagic rudd *Scardinius erythroptalmus* and piscivorous perch *Perca fluviatilis* and pike *Esox lucius*. Only two species were recorded from the Coolagh Lakes: roach *Rutilus rutilus* that are considered an invasive fish species, and perch. Only one specimen of European eel was also captured from the Coolagh Lakes despite connection to the River Corrib. No salmonids were recorded during the survey.

A summary of the numbers of fish captured and of the length frequency distributions for each species are summarised in Table 3.4 and Figures 3.12 and 3.13 below.

Table 3.5. Fish species recorded during fyke net surveys at Ballindooley Lough and Coolagh Lakes

Scientific Name	Common Name	Ballindooley Lough	Coolagh Lakes
<i>Esox Lucius</i>	Pike	3	
<i>Perca fluviatilis</i>	Perch	11	13
<i>Rutilus rutilus</i>	Roach		6
<i>Scardinius erythroptalmus</i>	Rudd	3	
<i>Tinca tinca</i>	Tench	6	
<i>Anguilla Anguilla</i>	Eel		1




Figure 3.12. Length Frequency distribution of fish species captured at Ballindooley Lough




Figure 3.13. Length Frequency distribution of fish species captured at Coolagh Lakes



Plate 3.14. Juvenile Perch *Perca fluviatilis* captured in the Coolagh Loughs



Plate 3.15. Roach *Rutilus rutilus* captured in the Coolagh Loughs



Plate 3.16. Tench *Tinca tinca* captured in Ballindooley Lough



Plate 3.17. Adult Perch captured in Ballindooley Lough



Plate 3.18. Rudd *Scardinius erythrophthalmus* captured in Ballindooley Lough



Plate 3.19. Pike *Esox lucius* captured in Ballindooley Lake

Fisheries Habitat Evaluation

Watercourses containing good salmonid habitat can be considered of at least high value local ecological importance (NRA, 2009). While some of the habitats contained salmonid habitat (e.g. the Tonabrocky Stream and the Bearna Stream) others had poorer quality fisheries habitat (e.g. Terryland River). In some instances, where salmonid habitat was not present or poor, habitat supporting European eel and or estuarine fish species was present. Where shallow stream gradients permitted (e.g. lower reaches of the Knocknacragh and Trusky Streams), good quality transitional nursery habitat for estuarine fish species was present. The evaluation of the stream sites and of the two lake habitats surveyed are appraised and summarised below.

Table 3.6. Fisheries Evaluation of Watercourses surveyed as part of the proposed road development

Watercourse Name	Characteristics (upper to middle reaches i.e. overlapping road footprint)	Characteristics (lower reaches)	Fisheries Notes	Evaluation
Sruthán na Libeirtí	The river is shallow (<0.3m) deep and seasonal, typically <0.5m wide. Heavily encroached by bracken, bramble and gorse scrub. Of no fisheries value in upper reaches. Upper reaches dried up during 2013-2014 hot summers.	Swift flowing with a depth of 0.5m. Historically straightened in middle reaches but retaining some good pool, riffle and glide sequences. Gobble & gravel beds and patches of soft sediment present.	The lower reaches have some moderate quality salmonid and European eel habitat. Only European eel present in small numbers.	Of local importance (lower value) for European eel. No salmonids present.
Trusky Stream	Shallow & small river in upper reaches (<1m wide & 0.2m deep) with some moderate quality salmonid habitat in upper reaches given riffle, glide and pool sequences. However, despite some salmonid habitat being present no salmonids were recorded.	The lower reaches of the Trusky Stream were wider (circa 3m) and deeper (0.5m) than the upstream habitat, given the confluence of three small tributaries. The habitat was dominated by faster flowing glide which was considered tidal during spring tides.	Lower reaches of some importance to brown trout and of high importance to European eel elver and juvenile flounder as a nursery habitat. Some spawning habitat for trout exists in the lower sections of the stream but the spawning areas are limited and the trout population size is considered small as a consequence.	Of local importance (higher value) for salmonids, European eel & as a nursery for flounder
Tonabrocky Stream	The upper reaches of the Tonabrocky Stream	The Lower reaches of the Tonabrocky	Upper reaches seasonal but moving	Of local importance (higher

Watercourse Name	Characteristics (upper to middle reaches i.e. overlapping road footprint)	Characteristics (lower reaches)	Fisheries Notes	Evaluation
	agglomerate were not considered of fisheries value. However moving downstream of the road alignment the habitat improves considerably with sequences of riffle, glide & localised pool. The gravel base of the stream also provides some spawning opportunity for trout. Small numbers of trout recorded upstream in the poorer quality habitat.	Stream have good quality salmonid habitat (both juvenile & adults). Some good quality spawning habitat present and localised holding pools for adult fish. Overall the stream increases considerably in size moving downstream and as such becomes a better salmonid habitat. In the lower reaches the stream had very swift flowing water with well defined riffle, pool and glide sequences.	downstream the habitat becomes an important salmonid river.	value) for brown trout
Bearna Stream	The Bearna Stream tributary of the Tonabrocky Stream is an excellent salmonid habitat throughout its upper reaches and considered the highest quality salmonid habitat of all of the tributaries surveyed along the corridor of the proposed road development.	Lower reaches contain excellent salmonid spawning and nursery habitat as do the upper reaches.	The Bearna Stream was considered an excellent nursery salmonid stream with good numbers of juvenile brown trout and small numbers of European eel. The habitat has excellent quality spawning habitat that was widespread.	Of local importance (higher value) for brown trout & European eel
Knocknacarragh Stream	The upper reaches of Knocknacarragh Stream were largely seasonal grassy ditches with the exception of one small section of channel downstream of Ballagh that retained some natural characteristics. Overall of no fisheries value.	The lower reaches of the Knocknacarragh Stream are heavily culverted. However a short section of open channel downstream of the R336 provides important habitat for juvenile estuarine fish and European eel.	Upper reaches of no fisheries value but lower reaches near estuary are of importance to as a transitional nursery habitat for estuarine fish & European eel.	Of local importance (higher value) for European eel & as a nursery for estuarine fish (sand goby, flounder & grey mullet)
Terryland River	The Terryland River in its upper	The lower reaches of the	Small numbers of European eel	Of local importance

Watercourse Name	Characteristics (upper to middle reaches i.e. overlapping road footprint)	Characteristics (lower reaches)	Fisheries Notes	Evaluation
	reaches is a heavily modified and polluted stream habitat of limited fisheries value.	Terryland River continue to be impacted by urban pollution and are of limited fisheries value.	recorded during the survey but otherwise of limited fisheries value.	(lower value) for European eel.
Ballindooley Lough	Ballindooley Lough is considered an excellent coarse fishery, but not of importance as a salmonid fishery. It has very clean water and has had low levels of human impact.		Good numbers of tench, pike, rudd and perch recorded indicating the lake is a coarse fishery.	Of local importance (higher value) for coarse fish species.
Coolagh Lakes	The Coolagh Lakes are of some value to coarse fish but are not considered of importance for salmonids. They are deep and steep shelving lakes with cold water fed by springs and the River Corrib.		Despite connection to the River Corrib of limited or no value to salmonids. Only coarse fish recorded during survey (i.e. Perch & Roach) & one European eel	Of local importance (lower value) for coarse fish species & European eel.

5. Discussion

Currently, the biological water quality of streams along the proposed road development are not achieving target good status (Q4), with the exception of the Tonabrocky Stream and its tributary the Bearna Stream. Rivers with good status typically are better quality fisheries. The baseline fisheries habitat and stock compositions are discussed below.

River Habitats

Salmonids

Atlantic salmon *Salmo salar* and sea trout *Salmo trutta* (migratory form) were not recorded present during the surveys, likely because the watercourses surveyed were small and shallow. Consequentially, accessible good quality spawning habitat for migratory salmonids was limited. Brown trout *Salmo trutta* were however recorded in good numbers in the Bearna Stream (site 3B), while small populations were also found present in the Tonabrocky Stream (site 4A) and in the lower Trusky Stream (site 2A). Given that these sites contain wild brown trout populations all efforts should be made to prevent or minimise impacts to river substrata, river profile and water quality. Wild Irish Brown trout populations are considered to be genetically diverse with numerous strains (Taggart et al. 1981; Ferguson, 2006) and, thus, are important for the wider conservation and management of the species in Europe.

The Trusky Stream may be considered the most vulnerable salmonid bearing stream overlapping the proposed road development given the low population size of trout in the stream. This reflected by the small numbers captured ($n=2$) and because of the very limited spawning, nursery and holding habitat present (i.e. restricted to 200m in lower reaches). As such every effort should be made to ensure that potential impacts from upstream at the proposed road development crossing do not impact the downstream habitat.

The middle-upper reaches of the Tonabrocky Stream also had low densities of trout. However, given that abundant higher quality salmonid bearing habitat was present downstream, it would have better prospects for recovery should water quality impacts damage the fishery. The Tonabrocky Streams major tributary, the Bearna Stream contained the highest quality salmonid habitat of all the stream sites surveyed. It had excellent nursery, spawning and holding habitat that was reflected by the healthy population structure comprising good densities of juvenile and adult trout. Given that both the Tonabrocky Stream and the Bearna Stream had good quality salmonid habitat it is important to prevent impacts to the existing excellent instream spawning habitat, good sinuous river profile and water quality. With regards the Bearna Stream where excellent salmonid habitat exists above the proposed road development crossing it is especially important to facilitate fish passage at the crossing.

European Eel

The critically endangered European eel *Anguilla anguilla* (Freyhoff & Kottelat, 2010) are considered to be the most threatened fish species in Ireland as indicated from a recent red listed publication on Irish Fish (King et al. 2011).

The European eel has protective status under the European Eel Regulation EC No. 1100/2007 to facilitate the recovery of the eel stocks since the large decline in the 1980's. Of the stream sites surveyed, the lower transitional reaches of the Trusky and Knocknacarragh Streams were good eel nurseries. Silver migratory adult eel were also recorded in the lower reaches of the Sruthán na Libeirtí Stream despite the upper reaches being devoid of fish. This exemplifies that the lower reaches of many of the stream catchments were of some value to European eel, even in the instances where the middle and upper reaches were of no fisheries value (e.g. Sruthán na Libeirtí, Trusky & Knocknacarragh). European eel were also recorded at low densities in the Bearna Stream and the Terryland River. Given that water borne pollutants can travel from the upper reaches of catchments to downstream, European eel nursery habitat within these reaches are especially important to protect (e.g. Trusky Stream). Similarly, where small and cryptic residual populations of eel persist in otherwise seasonal catchments, it is very important to prevent impacts to these fisheries. Measures would include prevention of downstream pollution, ensuring that downstream river profiles are not impacted, maintaining pockets of pool habitat and preserving fish passability. Recent evidence suggests that large numbers of juvenile 'glass' eels are returning to European rivers (Jackoby & Gollock, 2014) meaning it is important to capitalise on higher returning numbers by protecting both juvenile and adult eel habitat. This will help ensure numbers increase into the future given the large historical declines.

Lamprey species

According to Igoe et al. (2004) both sea lamprey *Petromyzon marinus* and brook lamprey *Lampetra planeri* are present in the River Corrib catchment (hydrometric areas 29 & 30; see Figure 2.1 for hydrometric divisions). Interestingly, no lamprey species have been recorded in hydrometric area 31 (Kelly & King, 2001) in which the majority of the surveyed stream sites are located. Detailed lamprey distribution surveys conducted by O' Connor (2007) indicated that lamprey distribution was patchy in the wider Corrib catchment and largely restricted to brook lamprey. While sea lamprey species are known to occur in the River Corrib they are now thought to be restricted below the Salmon Weir constructed in the 1960's (O' Connor, 2007).

Many of the watercourses surveyed for the proposed road development had compacted gravels and limited fine sediment deposition that are not conducive to larval lamprey settlement (e.g. Sruthán na Libeirtí and Trusky). Some localised areas of soft sediment were present within the Bearna Stream and the Tonabrocky Stream but more extensive areas of fines present in the Terryland River. Nonetheless, despite the presence of some suitable lamprey spawning and burrowing habitat, no larval lamprey were recorded at any of the survey sites. It must be noted, though, that this survey focused on small electro-fishing footprints area (typically between 25m² and 100m²). Therefore, the presence of larval lamprey further downstream of the proposed road development is considered possible in the Bearna Stream, the Tonabrocky Stream and Terryland River, albeit the noted suitable habitat was limited. This was considered following downstream fisheries habitat appraisals that identified localised larval lamprey habitat downstream of the electro-fishing areas.

The other stream habitats contained sub-optimal compacted cobble-gravel strewn streambed habitat that was contained in open, comparatively high-velocity streams (i.e. in the less seasonal tributaries). These included the Sruthán na Libeirtí, Trusky and Knocknacragh Streams.

In summary, potential for lamprey to occur downstream of electro-fished sites existed only in the Tonabrocky, the Bearna and Terryland catchments. While this may indicate the possible presence of low densities of juvenile lamprey downstream of the proposed road development, the Tonabrocky Stream and the Bearna Stream are located in hydrometric area 31 where no lamprey are known to occur (Kelly & King, 2001). While lamprey species are known to be present in the River Corrib to which the Terryland River is a tributary, gross pollution (i.e. Q2-3) was recorded in the Terryland River meaning the species is unlikely to persist as its tolerances typically relate to Q3 rivers and above.

Potential sea lamprey spawning areas downstream of the proposed road development are very unlikely. Sea lamprey typically utilise similar (or even the same) spawning areas to Atlantic salmon; spawning in coarse gravel, pebbles and sand, where the diameter of the gravel can vary from 1–11cm, the overlying water column has a depth of 40–60cm (Igoe *et al.*, 2004) and which are frequently found at the tail end of pools or conversely the upstream ends of rapids and riffles in relatively strong currents of up to 1–2ms⁻¹ (APEM, 2004). The water depths recorded in the surveyed streams with the exception of the Terryland River would unlikely support the species. The Terryland River itself was heavily polluted and primarily comprised soft silt channel bed and is not considered suitable for the sea lamprey adults. Furthermore, sea lamprey are thought to be restricted below the Galway Salmon Weir because it acts as a barrier to migration (O' Connor, 2007) and therefore they are unlikely to be able to access the Terryland River.

Estuarine Fish

Where the gradients of the lower reaches of stream habitat adjoining estuaries facilitates the deposition of gravels and fines they can be important fish nurseries for estuarine species. Two stream sites contained such habitat, i.e. the Trusky Stream & the Knocknacarragh Stream. Both sites had moderate flow rates and mixed sediment substrata comprising cobbles, gravels and coarse sand. The lower Trusky Stream was an excellent nursery for European eel elvers (young eel), flounder and three-spine stickleback despite poor fisheries habitat upstream. While the Knocknacarragh Stream contained similar habitat it was more saline and had the highest diversity of fish species of all the stream sites surveyed. Grey mullet *Chelon labrosus*, sand goby *Pomatoschistus minutus*, flounder *Platichthys flesus*, three spined stickleback *Gasterosteus aculeatus* and European eel were recorded as present. The presence of good transitional nursery habitat where stream sites discharge into estuaries exemplified the importance of stream catchments longitudinally as far downstream as the estuarine reaches. This is especially important when seemingly poor upstream and middle reach habitat exists, meaning the lower reaches can be overlooked, if one were to consider the upper catchments as an indicator of the importance of lower catchment alone.

Lake Habitats

Ballindooley Lough

Ballindooley Lough was considered an excellent mixed stock coarse fishery based on the findings of the survey. Rudd *Scardinius erythrophthalmus* were detected at low densities in one fyke net in the windward and shallow north basin of Ballindooley Lough. Here the expansive beds of *Chara spp.* and *Utricularia sp.* vegetation provide refugia for rudd and grazing opportunities. Furthermore, rudd, with their characteristic upturned mouth, feed on emerging aquatic insects that attach to submerged vegetation. As such, rudd as a species requires reeded littorals for cover and feeding. In contrast perch were detected in the deeper water, where the younger year classes would graze on plankton. Where rudd and perch co-exist in a medium sized waterbody i.e. 30 acres the later species tends to be numerically more abundant given they are superior planktivores in open water, in addition to becoming piscivorous as adults. Rudd in contrast to perch, have a herbivorous component to their diet eating Chara vegetation in addition to feeding aerial aquatic insects and zooplankton. As such in larger watercourses they tend to be restricted to shallower weeded bays (Kennedy & Fitzmaurice, 1973) as was found during the current survey.

Perch were recorded at moderate densities and in three fyke nets and appeared to be very numerous in the open and deeper water of the lake based on high resolution transducer readings. The absence of roach detected during the survey would indicate that the open water pelagic fish are likely to be perch rather than other species. Perch are considered a non-native fish species in Ireland (King et al., 2011).

Pike were detected in two fyke nets. The species also appeared in the deeper open water on transducer readings below perch shoals. Pike are a top predator and are obligate piscivores, but also eat small amphibians, mammals and waterfowl. They are nonetheless considered an important species for recreational angling and for the sustenance of balanced coarse fisheries (Arlinghaus et al., 2010). While pike were considered non-native historically more recent evidence suggests that early colonisation of the species may have been independent of humans (Pedreschi et al., 2013).

Tench were detected at moderate densities during the survey being recorded in 4 fyke nets on the western shore. The abundant beds of Chara vegetation and sheltered nature of the lake with limited direct cooling from rivers created a very good quality tench habitat in Ballindooley Lough. The species can thrive in small and medium sized waterbodies where marginal vegetated lake shelves in higher alkalinity lakes provide rich invertebrate feeding. They typically graze molluscs attached to Chara beds and rarely leave the seclusion of these habitats. As with pike, perch and rudd they are an important recreational angling quarry.

The basinal characteristics of Ballindooley Lough facilitate a habitat supporting number of different coarse fish species with different biological requirements. The open water provides good habitat for perch and pike with the weedy margins providing good habitat for rudd and tench.

Weeded margins also provide habitat for pike which use these areas for ambush predation. Ballindooley Lough is not considered of value as a game fishery (i.e. brown trout) as no salmonids were recorded during the survey. As it has no direct connection to adjoining rivers it is neither of importance to anadromous or catadromous fish. This conclusion is drawn based on the absence of both salmonids and European eel during the survey.

Overall Ballindooley Lough is considered a very high quality mixed coarse fishery. In this fashion it is a similar fishery to the nearby Ballyquirke and Ross Loughs to the south west of Lough Corrib, as both fisheries are considered also important as recreational coarse fisheries. These Lakes contrast to the internationally important game fisheries of Lough Corrib and Lough Mask that are famous for the brown trout and the ferox subspecies they support.

Coolagh Lakes

The Coolagh lakes are reed fringed and up to 18m deep as recorded on high resolution transducers. The two basins are connected by a narrow reed fringed channel with the northern basin being slightly shallower (12m) and spring fed. The lakes are largely inaccessible from the shoreline and were accessed by boat from a connecting channel via the River Corrib. Three species of fish were recorded at the Coolagh Lakes, namely roach, perch and European eel. Roach are classified as a 'non-native, non-benign' invasive species in Irish waters (Stokes *et al.*, 2004; King *et al.*, 2011) and is placed under restrictions according to Articles 49 and 50 of the S.I. 477 of the European Communities (Birds and Natural Habitats) Regulations 2011. Their environmental plasticity and ability to compete with other native fish species, such as Atlantic salmon and brown trout, for food means that they can have negative impacts on native fish populations. Unfortunately, roach are now widespread in the Corrib catchment colonising through the interconnecting channels and lakes. Given that the Coolagh Lakes are connected to the River Corrib, roach would have naturally colonised from this point. Interestingly despite this connection to the river no trout were recorded and only one specimen of European eel was found present. Overall the Coolagh Lakes are not considered of high fisheries value, rather of local importance as a coarse fishery.

7. References

- APEM (2001). Standardised sampling strategies and methodologies for condition assessment within SAC rivers for sea, river and brook lamprey and bullhead – Phase I Report. English Nature, Peterborough. 48 pp. Atherton, I., Bosanquet, S. & Lawley, M. (Eds.) (2010) Mosses & Liverworts of Great Britain & Ireland. British Bryological Society.
- Arlinghaus, R., Matsumura, S. & U. Dieckmann (2010) The conservation and fishery benefits of protecting large pike (*Esox lucius* L.) by harvest regulations in recreational fishing. *Biological Conservation* 143, 1444–1459
- Carle, F. L., and M. R. Strub (1978). A new method for estimating population size from removal data. *Biometrics* 34: 621-630.
- Crozier, W.W. and Kennedy, G.J.A. (1994). Application of semi-quantitative electro-fishing to juvenile salmonid stock surveys. *Journal of Fish Biology* 45: 159-164.
- Ferguson, A. (2006) Genetic differences among brown trout, *Salmo trutta* and their importance for the conservation and management of the species. *Freshwater Biology* 21(1): 35-46.
- Freyhof, J. & Kottelat, M. (2010). *Anguilla anguilla*. In: IUCN 2013. IUCN Red List of Threatened Species. Version 2013.2. <www.iucnredlist.org>. Downloaded on 1st November 2015.
- Harvey, J. and Cowx, I. (2003). Monitoring the River, Brook and Sea Lamprey *Lampetra fluviatilis*, *Lampetra planeri* and *Petromyzon marinus*. Conserving Natura 2000 Rivers, Monitoring Series No. 5. English Nature, Peterborough.
- Igoe, F., Quigley, D.T.G., Marnell, F., Meskell, E., O'Connor, W and Byrne, C. (2004). The sea lamprey *Petromyzon marinus* (L.), river lamprey *Lampetra fluviatilis* (L.), and brook lamprey *Lampetra planeri* (Bloch) in Ireland: General biology, ecology, distribution and status with recommendations for conservation. *Biology and Environment: Proceedings of the Royal Irish Academy* 104B: 43-56.
- Jacoby, D. & Gollock, M. (2014). *Anguilla anguilla*. In: IUCN 2014. The IUCN Red List of Threatened Species. Version 2014.1. <www.iucnredlist.org>. Downloaded on 3rd November 2014.
- Kelly, F. L. and King, J. J. (2001). A review of the ecology and distribution of three lamprey species, *Lampetra fluviatilis* (L.), *Lampetra planeri* (Bloch) and *Petromyzon marinus* (L.): a context for conservation and biodiversity considerations in Ireland. *Biology and Environment: Proceedings of the Royal Irish Academy* 101B: 165-185.
- King, J.J., Lehane, B.M., Wightman, G.D., Dooley, R. and Gilligan, N. (2011). Development and implementation of environmental protocols in river 2. Biogeographical Or Marine Level 12/09/2013 15:59:56 Page 1 of 5 Report on the main results of the surveillance under article 11 for annex II, IV and V species (Annex B) maintenance in Ireland. *Water and Environment Journal* 25: 422 - 428.
- Lockwood, Roger N. and J. C. Schneider (2000). Stream fish population estimates by mark and recapture and depletion methods. Chapter 7 in Schneider, James C. (ed.) 2000. Manual of fisheries survey methods II: with periodic updates. Michigan Department of Natural Resources, Fisheries Special Report 25, Ann Arbor.

- Maitland PS (2003). Ecology of river, brook and sea lamprey. Conserving Natura 2000 Rivers Ecology Series No. 4: English Nature, Peterborough
- Niven, A.J. & McCauley, M. (2013) Lamprey Baseline Survey No2: River Faughan and Tributaries SAC. Loughs Agency, 22, Victoria Road, Derry
- O Grady, M.F. (2006). Channels and Challenges. Enhancing Salmonid Rivers. Irish Freshwater Fisheries Ecology and Management Series: Number 4, Central Fisheries Board, Dublin, Ireland.
- O'Connor, W. (2007) A Survey of Juvenile Lamprey Populations in the Corrib and Suir Catchments. Irish Wildlife Manuals No. 26. National Parks and Wildlife Service, Department of Environment, Heritage and Local Government, Dublin, Ireland.
- Pedreschi, D., Kelly-Quinn, M., Caffrey, J., O'Grady, M. & Mariani, S. (2013) Genetic structure of pike (*Esox lucius*) reveals a complex and previously unrecognized colonization history of Ireland. *Journal of Biogeography* 1-13. <http://doi:10.1111/jbi.12220>
- Stokes, K., O'Neill, K. & McDonald, R.A. (2004) Invasive species in Ireland. Unpublished report to Environment & Heritage Service and National Parks & Wildlife Service. Queens, Queens University Belfast, Belfast.
- Taggart, J, Ferguson, A., Mason, F.M. (1981) Genetic variation in Irish populations of brown trout (*Salmo Trutta*); electrophoretic analysis of allonyms. *Journal of Physiology & Biochemistry part B: Comparative Biochemistry* 69(3): 393-412.
- Thompson, K., Bronston, J.K. & Lozier, C.W. (2010). Best management practices to minimize adverse effects to Pacific lamprey (*Entosphenus tridentatus*). Columbia River Basin. 25pp.
- Toner, P., Bowman, J., Clabby, K., Lucey, J., McGarrigle, M., Concannon, C., Clenaghan, C., Cunningham, P., Delaney, J., O'Boyle, S., McCarthaigh, M., Craig, M. & Quinn, R. (2005) Water Quality in Ireland, 2001–2003. Environmental Protection Agency, Co. Wexford, Ireland.
- White, G. C., D. R. Anderson, K. P. Burnham, and D. L. Otis (1982). Capture-recapture and removal methods for sampling closed populations. Los Alamos: Los Alamos National Laboratory.
- Zalewski, M. & I.G. Cowx (1990) Factors affecting the efficiency of electric fishing. In: *Fishing with electricity: Applications in freshwater fisheries management*. Cowx, I.G. & Lamarque, P. (eds.): 89-112. Fishing News Books, Oxford.



Appendix A – Electro-fishing License



**Roinn Cumarsáide,
Fuinnimh & Acmhainní Nádurtha**
Department of Communications,
Energy & Natural Resources

**CERTIFICATE OF AUTHORISATION UNDER SECTION 14 OF THE FISHERIES
(CONSOLIDATION) ACT, 1959 AS SUBSTITUTED BY SECTION 4 OF THE
FISHERIES (AMENDMENT) ACT, 1962.**

The Minister for Communications, Energy and Natural Resources in exercise of the powers conferred on him by Section 14 of the Fisheries (Consolidation) Act, 1959 as substituted by Section 4 of the Fisheries (Amendment) Act, 1962 hereby authorises:

Mr Ross Macklin, Triturus Environmental Services, 42 Norwood Court,

Rochestown, Cork City. and or person(s) nominated by him to undertake an electro fishing surveys in relation to proposed N6 Galway City Transport Project.

The works involve the undertaking of quantitative electro-fishing surveys of small rivers/ streams (n=7) along the alignment of the proposed N6 Galway City Transport Project and the quantitative assessment of the fisheries status of two lake systems that are connected hydrologically to the scheme, by means of fyke netting.

The following locations have been proposed for the survey: 1. Liberty (West of River Corrib); 2. Trusky (West of River Corrib); 3. Bearnna (West of River Corrib); 4. Rusheen Bay (West of River Corrib); 5. Coolagh lakes stream (East of River Corrib); 6. Terryland Stream (East of River Corrib); 7. Merlin Park Stream (East of River Corrib); 8. Coolagh Lakes (East of River Corrib); and 9. Ballindooly Lough (East of River Corrib). Electro-fishing will be undertaken in locations 1-7 and fyke netting will be undertaken in locations 8 and 9 (site details and map are included in the original application).



This authorisation is granted subject to the following conditions:

- 1. This authorisation shall not confer on the holder thereof, independently of the conditions therein;**
 - (a) any rights or title which the holder would not have had if this Authorisation had not been given, or;**
 - (b) any authority in any way to interfere with or infringe the lawful rights of any other person.**
- 2. This authorisation is issued to and valid for use by Bill Quirke and or person(s) nominated by him.**
- 3. This authorisation is valid until 30 September 2015.**
- 4. Inland Fisheries Ireland Galway (IFI) shall be notified at least 5 working days in advance of the proposed commencement of the electro-fishing operations. Please contact IFI Director John Conneely / Pat Gorman, IFI Fisheries Inspector. The holder of this authorisation shall comply with any instructions given to them in relation to fishing operations.**
- 5. IFI recommends that the applicant seeks permission from fishery owners and informs local angling clubs of their plans for the surveys where relevant. The applicant must also seek permission from landowners to cross land, where relevant.**
- 6. The electrofishing operation must be carried out during suitable weather and flow conditions.**



- 7. Electro-fishing should, if possible, be carried out by the September 30th when juvenile salmonids (if present) are of a sufficiently large size to be caught by electro-fishing, to minimize damage and for them to be distinguished from similar species (CEN, 2001 and CFB/IFI 'Electric Fishing in Wadeable reaches' manual.**

- 8. Fyke netting operations should be carried out by the 30th September 2015.**

- 9. The applicant is fully aware of biosecurity concerns. IFI insists upon strict adherence to the Biosecurity Protocol for Field Survey Work, whereby equipment must be disinfected prior to and after use to prevent the spread of disease, parasites or invasive species**
[\(<http://www.fisheriesireland.ie/Biosecurity/biosecurity-protocol-for-field-survey-work.html>\)](http://www.fisheriesireland.ie/Biosecurity/biosecurity-protocol-for-field-survey-work.html) (and as directed by an officer of IFI).

- 10. All equipment must be available for inspection by an IFI officer during the survey.**

- 11. The fishing gear when not in use shall be kept in a secure place known to an Officer of the Western River Basin District of IFI and the local Garda Siochana.**

- 12. IFI request that any crayfish captured in the survey be measured (carapace length in mm is standard). This information should also be included on the IFI reporting template.**



**Roinn Cumarsáide,
Fuinnimh & Acmhainní Náúúrtha**
Department of Communications,
Energy & Natural Resources

- 13. Any fish captured shall be carefully handled and returned alive to the water from which they are taken, following the gathering of data. No fish of any species should be sacrificed during the appraisal, except in circumstances where tissue/body burden analysis is to be carried out. IFI Galway shall be informed of any fish mortalities immediately after the survey. Details including the County, Site number, River Name, Townland, Irish Grid Reference and the species.**

- 14. When doing anything pursuant to this authorisation, the holder shall, if requested by any person affected, produce this authorisation to that person.**

- 15. The survey report including the survey quantitative data obtained in the appraisal shall be forwarded electronically in the standard IFI format to Sandra Doyle, Inland Fisheries Ireland, 3044 Lake Drive, Citywest Business Campus, Dublin 24, Sandra.doyle@fisheriesireland.ie within 30 days of completion of the survey. For ease of reference an electronic copy of IFI's standard template will be forwarded directly to the applicant. Report (hard and soft copies) to include mortality data and full account of qualitative/quantitative results. These data will not be made publically available, for a period of three years, without the permission of Triturus Environmental Services.**

- 16. Failure to comply with any of the conditions of this authorisation may result in revocation of this authorisation.**

- 17. The holder of this authorisation should be mindful of the potential occurrence of invasive alien species be recorded, IFI request that their location is recorded and submitted to IFI. It is likely that Japanese knotweed will be encountered in or around the survey locations.**



**Roinn Cumarsáide,
Fuinnimh & Acmhainní Nádurtha**
Department of Communications,
Energy & Natural Resources

18. The holder of this authorisation shall indemnify and keep indemnified the State, the Minister for Communications, Energy and Natural Resources and the Minister for Finance against any claims, arising in any manner whatsoever in connection with the user of the fishing gear or in the exercise of the permission hereby granted.

19. Notwithstanding the foregoing, this authorisation may be revoked or amended by the Minister for Communications, Energy and Natural Resources without the payment of compensation to the holder on giving one week's notice in writing to the holder if he considers it necessary in the public interest to do so.

Dated this 11 September 2015





For the Minister for Communications, Energy and Natural Resources.

Gerry Clerkin

An officer authorised on that behalf by the said Minister



Appendix B – Lake Profiles





Appendix C – Fyke Net Locations




Figure C1. Location of Fyke Netting Sites on Ballindooley Lough




Figure C2. Location of Fyke Netting Sites on the Coolagh Lakes